Model Question Paper

Applications of matrices and determinants - Part II

Business Maths	Reg.No. :			

I.Answer all the questions. II.Use Blue pen only. III.Question No 15 is compulsory

Time: 01:15:00 Hrs Total Marks: 85

 $5 \times 1 = 5$

1) The inverse of $\begin{pmatrix} 0 & 2 \\ 2 & 0 \end{pmatrix}$ is

(a)
$$\begin{pmatrix} 0 & \frac{1}{2} \\ 2 & 0 \end{pmatrix}$$
 (b) $\begin{pmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{pmatrix}$ (c) $\begin{pmatrix} 0 & -\frac{1}{2} \\ \frac{1}{2} & 1 \end{pmatrix}$ (d) $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$

(a)
$$\begin{pmatrix} 0 & \frac{1}{2} \\ 2 & 0 \end{pmatrix}$$
 (b) $\begin{pmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{pmatrix}$ (c) $\begin{pmatrix} 0 & -\frac{1}{2} \\ \frac{1}{2} & 1 \end{pmatrix}$ (d) $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$
2) If $A = \begin{pmatrix} 0.8 & 0.6 \\ -0.6 & 0.8 \end{pmatrix}$ then A^{-1} is

(a) $\begin{pmatrix} -0.8 & 0.6 \\ -0.6 & 0.8 \end{pmatrix}$ (b) $\begin{pmatrix} 0.8 & -0.6 \\ 0.6 & 0.8 \end{pmatrix}$ (c) $\begin{pmatrix} 0.8 & 0.6 \\ 0.6 & 0.8 \end{pmatrix}$ (d) $\begin{pmatrix} 0.2 & 0.4 \\ -0.4 & 0.2 \end{pmatrix}$

Section-A

For what value of k the matrix A, where $A=\begin{pmatrix} 2 & k \\ 3 & 5 \end{pmatrix}$ has no inverse?

(a)
$$\frac{3}{10}$$
 (b) $\frac{10}{3}$ (c) 3 (d) 10
4) If $A = \begin{bmatrix} 2 & 3 & 1 \\ 3 & 4 & 1 \\ 3 & 7 & 2 \end{bmatrix}$ then A^{-1} A is (a) 0 (b) A (c) I (d) A^2

5) The rank of an n x n matrix each of whose elements is 1 is

(a) 1 (b) 2 (c) n (d)
$$n^2$$

5 x 6 = 30 Section-B

Write the inverse of $A=\begin{pmatrix} 2 & 4 \\ -3 & 2 \end{pmatrix}$

Write the inverse of $A=\begin{bmatrix} 1&0&2\\1&0&2\\3&1&1\\2&1&2\end{bmatrix}$ Find the inverse of $A=\begin{bmatrix} 1&0&a\\0&1&b\\0&0&1\end{bmatrix}$ and verify that $AA^{-1}=I$.

If $A=\begin{bmatrix}a_1&0&0\\0&a_2&0\\0&0&a_3\end{bmatrix}$ and none of the a's are zero, find A^- If $A=\begin{bmatrix}-1&2&-2\\4&-3&4\\&&&1\end{bmatrix}$ show that the inverse of A is itself.

Section-C $5 \times 10 = 50$

- 11) There are three commodities X, Y and Z which are bought and sold by three dealers A, B and C. Dealer A purchases 2 units of X and 5 units of Z and sells 3 units of Y, dealer B purchases 5 units of X, 2 units of Y and sells 7 units of Z and dealer C purchases 3 units of Y, 1 unit of Z and sells 4 units of X. In the process A earns Rs.11 and C Rs.5 but B loses Rs.12. Find the price of each of the commodities X, Y and Z, by using determinants.
- 12) A company produces three products everyday. The total production on a certain day is 45 tons. It is found that the production of the third product exceeds the production of the first product by 8 tons while the total production of the first and third product is twice the production of second product. Determine the production level of each product by using Cramer's rule.
- 13) The data below are about an economy of two industries P and Q. The values are in millions of rupees.

Producer			User		Final Demand	Total Output	
Producer	Р	Q	Total Output				
Р	14	6	8	28			
Q	7	18	11	36			

Determine the outputs if the final demand changes to 20 for P and 30 for Q.

14) Suppose the inter-relationship between the production of two industries P and Q in a year (in lakhs of rupees) is

Producer	User		Final Damand	Tatal Outrot	
	Р	Q	Final Demand	Total Output	
Р	15	10	10	35	
Q	20	30	15	65	

Find the outputs when the final demand changes to (i) 12 for P and 18 for Q (ii) 8 for P and 12 for Q.

15) a) In an economy of two industries P and Q the following table gives the supply and demand positions in millions of rupees.

Ī	Producer	User		Final Demand	Total Output	
		Р	Q	Filial Dellialiu	Total Output	
ĺ	Р	16	20	4	40	
ĺ	Q	8	40	32	80	

Find the outputs when the final demand changes to 18 for P and 44 for Q.

(OR)

b) The data below are about an economy of two industries P and Q. The values are in crores of rupees.

Producer	User		Final Demand	Total Output	
Producer	Р	Q	rillat Dellialiu	Total Output	
Р	50	75	75	200	
Q	100	50	50	200	

Find the outputs when the final demand changes to 300 for P and 600 for Q.



